

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED FINAL/15 DEC 93 TO 14 DEC 94
----------------------------------	----------------	--


4. TITLE AND SUBTITLE RAREFIED GAS DYNAMICS AND MONTE CARLO METHODS	5. FUNDING NUMBERS
--	--------------------

6. AUTHOR(S) RUSSEL E. CAFLISCH	2304/BS F49620-94-1-0091
------------------------------------	-----------------------------

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) UNIVERSITY OF CALIFORNIA MATHEMATICS DEPARTMENT LOS ANGELES, CA 90024-1406	8. PERFORMING ORGANIZATION REPORT NUMBER AFOSR-TR-95-0430
--	---

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NM 110 DUNCAN AVE, SUITE B115 BOLLING AFB DC 20332-0001	10. SPONSORING/MONITORING AGENCY REPORT NUMBER F49620-94-1-0091
--	---

11. SUPPLEMENTARY NOTES



12a. DISTRIBUTION/AVAILABILITY STATEMENT
--

12b. DISTRIBUTION CODE

APPROVED FOR PUBLIC RELEASE: DISTRIBUTION IS UNLIMITED
--

13. ABSTRACT (Maximum 200 words)

The research supported by this grant was primarily on quasi-random sequences, which are a deterministic alternative to random or pseudo-random sequences, and their use in Monte Carlo methods. These methods are vastly superior to standard Monte Carlo, at least in principle, because the errors are of size $O(N^{-1})$ rather than the usual $O(N^{-1/2})$. On the other hand, this gain in accuracy can be lost if the domain of integration is of high dimension or the function to be integrated is not smooth. In joint work with Bradley Moskowitz, we succeeded in overcoming these limitations for a number of Monte Carlo methods, including the acceptance-rejection method, Feynman-Kac integrals and diffusion Monte Carlo.

DTIC QUALITY INSPECTED 3

14. SUBJECT TERMS		15. NUMBER OF PAGES
		16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED
20. LIMITATION OF ABSTRACT SAR(SAME AS REPORT)		

1995061901

Accession For)	
NTIS	CRA&I	<input checked="" type="checkbox"/>	
DTIC	TAB	<input type="checkbox"/>	
Unannounced		<input type="checkbox"/>	
Justification			
By			
Distribution /			
Availability Codes			
Dist	Avail and / or Special		
A-1			

FINAL TECHNICAL REPORT
AFOSR Grant 94-1-0091
Rarefied Gas Dynamics and Monte Carlo Methods

Russel E. Caflisch, PI
Mathematics Department, UCLA

May 19, 1995

The research supported by this grant was primarily on quasi-random sequences, which are a deterministic alternative to random or pseudo-random sequences, and their use in Monte Carlo methods. These methods are vastly superior to standard Monte Carlo, at least in principle, because the errors are of size $O(N^{-1})$ rather than the usual $O(N^{-1/2})$. On the other hand, this gain in accuracy can be lost if the domain of integration is of high dimension or the function to be integrated is not smooth. In joint work with Bradley Moskowitz, we succeeded in overcoming these limitations for a number of Monte Carlo methods, including the acceptance-rejection method, Feynman-Kac integrals and diffusion Monte Carlo.

In particular we developed the following new method that employ quasi-random sequences:

- (1) A smoothed acceptance-rejection method.
- (2) A rearrangement of the Feynman-Kac integral using the Brownian bridge.
- (3) A hybrid method for diffusion Monte Carlo, using pseudo-random to get a good approximation to the solution, then quasi-Monte Carlo to reduce the statistical error.

Funds from this grant were used to support Bradley Moskowitz as a postdoctoral researcher and a visitor, Giovanni Russo. The duration of this grant was one year.

Publications Supported by This Grant

1. B. Moskowitz, "Improved Stochastic Simulation Using Quasi-Monte Carlo: A Computational Study" UCLA CAM Report 94-40
2. B. Moskowitz, "Quasirandom Diffusion Monte Carlo" UCLA CAM Report 94-39
3. B. Moskowitz and R. Caflisch, "Smoothness and Dimension Reduction in Quasi-Monte Carlo Methods" UCLA CAM Report 94-17, to appear in J. Math. Comp. Modeling

Publications Supported by Previous AFOSR Grants that Appeared During this Period

1. W. Morokoff and R. Caflisch "A Quasi-Monte Carlo Approach to Particle Simulation of the Heat Equation" SIAM J. Num. Anal. 30 (1993) 1558-1573.
2. W. Morokoff and R. Caflisch "Quasi-Random Sequences and Their Discrepancies" SIAM J. Sci. Stat. Comp. 15 (1994) 1251-1279.